

Supporting Information

Hafnia-silica cryogels: solvent-assisted textural and catalytic control in the citronellal cyclization.


Ciril Jimeno,^{*,[a]} Jonathan Miras,^[b] and Jordi Esquena^[b]

[a] Department of Biological Chemistry and Molecular Modelling
Institute of Advanced Chemistry of Catalonia (IQAC-CSIC)
c/ Jordi Girona 18-26, E08034, Barcelona (Spain)
Fax: (+)34 93 2045904
E-mail: ciril.jimeno@iqac.csic.es


[b] Department of Chemical and Biomolecular Nanotechnology
Institute of Advanced Chemistry of Catalonia (IQAC-CSIC) and CIBER de Bioingeniería,
Biomateriales y Nanomedicina (CIBER-BBN)
c/ Jordi Girona 18-26, E08034, Barcelona (Spain)


1. Powder XRD of Hf-Si8-X	S2
2. FTIR spectra of Hf-Si8-X	S3
3. Sorption isotherms and pore size distributions for Hf-Si8-D, -E, and -F	S5
4. Extra SEM images of Hf-Si-A, -B, and -C	S8
5. Subtracted TGA of adsorbed pyridine on Hf-Si-A, -B, and -C	S10

1. POWDER XRD OF Hf-Si8-X.




2. FTIR SPECTRA OF Hf-Si8-X.






3. SORPTION ISOTHERMS AND PORE DISTRIBUTIONS FOR Hf-Si8-D, -E, AND -F.




Hf-Si8-D;



Hf-Si8-B;






Hf-Si8-C;





4. EXTRA SEM IMAGES OF Hf-Si8-A, -B, AND -C.




Hf-Si8-A;



Hf-Si8-B;



Hf-Si8-C;



5. SUBTRACTED TGA OF ADSORBED PYRIDINE ON Hf-Si-A, -B, AND -C

